

Companion Tests Provide Common Platform for Psychologists and Medical Practitioners in Evaluating Pain Patients

A CASE STUDY

Since its founding in 1995, The Pain Treatment Center at Rush Foundation Hospital has been committed to taking a multidisciplinary approach in treating pain patients. The clinic recognized early on the value of administering psychological assessments that would help the psychologists, physicians, and other members of the care team work together effectively and efficiently. After using a variety of available tools, the team has found that the BHI™ 2 (Battery for Health Improvement 2) test and the shorter version of this instrument, the BBHI™ 2 (Brief Battery for Health Improvement 2), are best suited to their needs.

The following article outlines the center's evolution as a multidisciplinary service and the benefits that the clinic's psychologists and medical practitioners have discovered in using the BHI 2 and BBHI 2 tests.

Developing a multidisciplinary team

When Rush Foundation Hospital in Meridian, Mississippi, hired pain specialist Eric Pearson, MD, to head up their anesthesiology department in 1995, one of his first priorities was to persuade the hospital to build a pain center. Since moving to Meridian in 1992, Dr. Pearson had become keenly aware of the lack of pain management services in the region.

The hospital readily agreed to Dr. Pearson's proposal. The initial plan called for Dr. Pearson to divide his time equally between the pain clinic and the anesthesiology department. The demand for pain management services in the region was so great that within two weeks after the Pain Treatment Center at Rush Foundation Hospital opened its doors in January of 1996, Dr. Pearson found himself working full time as director at the pain clinic.

From the onset, Dr. Pearson focused on building a multidisciplinary team to meet the biopsychosocial needs of pain patients. Early on, the clinic subcontracted with a group of psychologists in the area, and by 1998 it had staffed two additional pain specialist psychologists. Over the years, the clinic has steadily expanded the care team. The current group also includes two pain specialist physicians, two professional counselors, a nurse practitioner, a physician assistant, more than 30 nurses, and a physical therapist. In addition, the clinic makes referrals to outside services, including chiropractic and massage therapy.

The Pain Treatment Center at Rush Foundation Hospital

James Gilbert, PhD James McLaughlin, PhD Eric Pearson, MD

"Highly elevated scores on the BHI 2 test's Defensiveness, Substance Abuse, and Doctor Dissatisfaction scales help alert us that the patient is not a likely candidate for narcotics therapy."



James Gilbert, PhD, Coordinator of Behavioral Medicine at the clinic, explains why it is so important to use a multidisciplinary model in treating pain patients. "Many of our clients have already reached the chronic stage by the time they come to us. We offer different tools to help them manage their symptoms, but these symptoms are in the context of a life; patients must learn how to manage pain on a daily basis. Through a lifespan approach, we help patients rehabilitate physically and emotionally—and we help them set realistic expectations."

The clinic has built a broad referral radius, with the majority of referrals coming from physicians in the medical community. It serves many different types of pain patients, including individuals with failed back syndrome, complex regional pain syndrome, cancer pain, and neuropathic pain. About 20 percent of the clinic's clientele are workers' compensation cases.

Identifying the right tools

Over the years, the clinic has used various psychological assessments to identify those that are most relevant in addressing its patients' multifaceted issues. In 2002, Dr. Gilbert introduced the use of the BHI 2 test and its shorter version, the BBHI 2 test. Both assessments have since become a standard part of the clinic's protocol.

"The BBHI 2 and BHI 2 tests appealed to us because they are multiscale instruments that look at physical, emotional, and psychosocial aspects," says Dr. Gilbert. "They give us a better understanding of the patient's full pain experience, which goes far beyond just physical symptoms." Dr. Pearson notes that the tests provide valuable information on psychosocial factors commonly seen in pain patients, such as pain, somatic, and functional complaints—as well as psychological concerns such as depression, anxiety, and patient defensiveness.

"Another big plus of these tests is that they are normed on a physical injury population, making them more relevant to our patients than most of the other tests available, which are typically normed on a psychiatric population," says James McLaughlin, PhD, one of the clinic's psychologists. "In addition, the construction of the BBHI 2 and BHI 2 tests help reduce the risk of overpathologizing pain patients. For example, chronic pain patients may exhibit vegetative symptoms such as sleep irregularities and appetite disturbance that are commonly associated with depression, and yet these symptoms may be mainly related to the pain and the person may not be seriously depressed. The BBHI 2 and BHI 2 tests are designed to help sort out these confounding symptoms."

A two-step approach

Every patient at the clinic completes the BBHI 2 test during the initial visit. Dr. Gilbert uses the test results to assess to what degree the behavioral medicine

"The BBHI 2 and BHI 2 tests appealed to us because they are multiscale instruments that look at physical, emotional, and psychosocial aspects," says Dr. Gilbert. "They give us a better understanding of the patient's full pain experience, which goes far beyond just physical symptoms."

specialists should be involved in the patient's care. "Some people may only need periodic follow-up to monitor progress and compliance; others may benefit from a more thorough psychological evaluation and ongoing therapy," he says. "The BBHI 2 test helps us determine the suitable level of assistance."

If there are significant subscale elevations on the BBHI 2 test, the behavioral medicine specialists attempt to gain greater perspective on the patient's issues. They administer the BHI 2 test as part of a more extensive evaluation. The follow-up evaluation also may include either the MMPI®-2 (Minnesota Multiphasic Personality Inventory®-2) or the MBMD® (Millon® Behavioral Medicine Diagnostic) test, depending on the clinician's preference.

The clinic administers the BHI 2 test to assess all candidates for implantable devices such as drug infusion systems and spinal implants. "It is especially important to verify the candidacy of the patient when one is considering an expensive invasive surgery, such as a percutaneous discectomy, that insurers are very resistant to covering," says Dr. Pearson. "In this area, the BHI 2 test's Symptom Dependency scale has proven very useful. We have found it to be the highest predictor of whether or not the patient is going to respond to treatment."

The BHI 2 test also is administered to all candidates for chronic narcotics therapy. "It is not easy for us to identify patients at risk for addiction simply through a physical exam and clinical interview," says Dr. Pearson. "Highly elevated scores on the BHI 2 test's Defensiveness, Substance Abuse, and Doctor Dissatisfaction scales help alert us that the patient is not a likely candidate for narcotics therapy."

In addition, the clinic gives the BHI 2 test to all workers' compensation patients. "In these cases, it can be very difficult to establish whether the patient's pain prevents them from returning to work or whether there are secondary issues at play," says Dr. Pearson. "Physical examinations often produce inconsistent findings. The BHI 2 test provides an empirical measure to help accurately reflect the patient's true pain experience."

Special report features add value

A component of the BHI 2 report that Dr. Gilbert finds especially useful is the section on treatment recommendations. "This feature is very 'clinician-friendly' for the treating psychologist or counselor," says Dr. Gilbert. "It offers specific suggestions about issues to address in counseling that can help ensure medical intervention will be successful for the patient."

An aspect of the report that Dr. McLaughlin appreciates is the Pain Tolerance Index, which compares the patient's self-reported pain level to his/her self-reported coping ability. "With this index, we gain a very helpful snapshot of the patient's perception of his/her level of disability, especially when coupled with the Functional Complaints score," he says.

"We have found it [The BBHI 2 test's Symptom Dependency scale] to be the highest predictor of whether or not the patient is going to respond to treatment."

Use of companion tests creates efficiencies

The care team cites several advantages to including the BBHI 2 and the BHI 2 tests in one's protocol. "Whether you are a staff psychologist with an in-house multidisciplinary group, or you are working on a referral basis with medical providers, the use of these companion tests can help ensure that you and the other members of the care team are speaking the same language in assessing the patient's needs," says Dr. Gilbert.

Because the tests derive from the same base and use the same terminology, administering both instruments also has helped the clinic clean up its data, says Dr. Pearson. "We are using fewer tools now to get the information we need," he comments.

Dr. McLaughlin points out an additional benefit to using both tests. "When we use the BHI 2 test to conduct a follow-up evaluation, we can conduct a more focused search; we are able to gain in-depth information on the same specific areas that were flagged by the BBHI 2 test. In addition, since the two tests measure the same core items, using both of them allows us to track the patient's status at two points in time—which helps inform our initial decisions about the patient's care."

Taking the vital signs

In their ongoing commitment to finding the most effective approach to patient care, the clinic has discovered that the BHI 2 and BBHI 2 tests are targeted to their needs. "These tests include specific scales that measure the emotional vital signs of the chronic pain patients we serve," says Dr. Gilbert. "By helping us to better understand the individual experience of pain, the tests help us to structure a more efficient and successful treatment plan for each patient."

"By helping us to better understand the individual experience of pain, the tests help us to structure a more efficient and successful treatment plan for each patient."

BBHI™ 2: Get the Full Picture of Your Pain Patients

James Gilbert, PhD, is Coordinator of Behavioral Medicine for the Pain Treatment Center at Rush Foundation Hospital in Meridian, Mississippi. A licensed professional counselor, Dr. Gilbert received his doctoral degree in health psychology from Capella University and his MSCE in counseling psychology from the University of West Alabama. His special interests include the psychological aspects of pain treatment, stress management, and working with public safety professionals.

James McLaughlin, PhD, recently assumed the position of Senior Mental Health Worker at the new Pain Management Center of Baylor All Saints Medical Center in Fort Worth, Texas. Previously, Dr. McLaughlin served as a staff psychologist for the Pain Treatment Center at Rush Foundation Hospital in Meridian, Mississippi. His role there included individual and group psychotherapy, behavioral medicine management, and clinical research. Dr. McLaughlin earned his doctorate in counseling psychology from the University of Southern Mississippi and is a diplomate of the American Academy of Pain Management.

Eric Pearson, MD, is board-certified in anesthesiology with a subspecialty certificate in pain management. He is founder and director of the Pain Treatment Center at Rush Foundation Hospital in Meridian, Mississippi. Dr. Pearson also founded the Department of Anesthesiology for the Rush Medical Group in 1996 and served as that department's director from 1996 to 2000. He received his doctoral degree from the Medical University of South Carolina and is a diplomate of the American Academy of Pain Management.

This article was originally published by Pearson in 2005.

866.335.8418 | PearsonClinical.ca

Copyright © 2018 Pearson Education. All rights reserved. Pearson, BHI, and BBHI are trademarks, in the US and/or other countries, of Pearson Education, Inc. or its affiliates. The following are registered trademarks of the Regents of the University of Minnesota: MMPI, MMPI-2-RF, MMPI-A-RF, Minnesota Multiphasic Personality Inventory, and Minnesota Multiphasic Personality Inventory-2-Restructured Form. The following are unregistered, common law trademarks of the University of Minnesota: MMPI-2, MMPI-A, Minnesota Multiphasic Personality Inventory-2, Minnesota Multiphasic Personality Inventory-Adolescent, Minnesota Multiphasic Personality Inventory-Adolescent-Restructured Form, and The Minnesota Report. MBMD and Millon are registered trademarks of DICANDRIEN, Inc. CLINA15777-10475B EL 2/18-5/21-ca

