

K-CAT[®]

Powered by **ADAPTIVE TESTING**
TECHNOLOGIES

K-CAT[®]

Multirater Report (Parent & Child)

Name: Johnny Sample

Birth date: 06/04/2013

Gender: Male

Administration language: English (US)

Test date: 09/17/2026

Report date: 09/23/2026

Age at testing: 13

Examiner name:

Timeframe: Past 2 weeks

Copyright © 2026 NCS Pearson, Inc. and Adaptive Testing Technologies. All rights reserved.

Warning: This report contains copyrighted material and trade secrets. The qualified licensee may excerpt portions of this output report, limited to the minimum text necessary to accurately describe their significant core conclusions, for incorporation into a written evaluation of the examinee, in accordance with their profession's citation standards, if any. No adaptations, translations, modifications, or special versions may be made of this report without prior written permission from Pearson.

Pearson and **Q-global** are trademarks, in the U.S. and/or other countries, of Pearson PLC or its affiliates. **K-CAT** is a trademark of Adaptive Testing Technologies.

[1.0 / RE1 / QG1]



This report is intended for use by qualified professionals trained in the use of psychological tests. Results should always be interpreted with consideration of clinical judgment, additional assessment data, and the psychosocial context. The following tables present the results of each module completed by the examinee. If selected, administered items and their ratings are included within the corresponding module section of the report. Additional guidance on interpreting each K-CAT® module is available in the K-CAT® Interpretive Guide in the Q-global® Resource Library.

Interpreting the K-CAT®

Each K-CAT® module provides four key metrics:

- **Category:** A clinical classification (e.g., normal, mild, moderate, or severe) based on severity scores. These may also be labeled as low risk, intermediate risk, or high risk, depending on the module.
- **Severity:** A score on a 100-point scale indicating symptom severity, with higher scores reflecting greater severity.
- **Precision:** A measure of the degree of uncertainty in the severity estimate. Adaptive testing allows the precision to be fixed at 5 points on a 100-point scale, where 0 would represent absolute certainty and 100 would represent absolute uncertainty.
- **Diagnostic Probability:** The probability that a child would receive a given diagnosis if they were evaluated using a full diagnostic interview.

The severity score and category indicate where the score falls in relation to the general population on which the module has been tested. The precision indicates how much certainty there is in the severity score. For example, a severity score of 75 with a precision of 5.0 indicates that the true score falls between 70 and 80. Repeated assessments would place the true score within 65 to 85, 95% of the time, forming a 95% confidence interval (i.e., 95% confidence interval = severity score plus/minus $1.965 \times \text{precision}$).

IMPORTANT: If the Suicide module was administered, carefully review the risk category and individual responses, as they provide critical information related to suicide risk that may require immediate attention or further evaluation.

Note. Duration is expressed in seconds in each table where it appears. If a duration is not shown for an item (-), the item was administered as part of another module and the response was used for the module in question without re-administration.

Test Results

Timeframe: Past 2 weeks

Results Summary

Module	Category		Severity		Precision	
	Child	Parent	Child	Parent	Child	Parent
Suicide	Moderate		47.2		4.6	
Anxiety	Mild	Normal	34.0	18.1	6.4	5.0
Mania	Mild	Moderate	32.1	60.3	5.5	4.6
Oppositional defiant disorder (ODD)	Severe	Severe	66.7	80.1	4.3	4.2
Attention deficit/hyperactivity disorder (ADHD)	Moderate	Moderate	57.9	62.0	6.0	3.4
Depression	Mild	Mild	48.9	40.4	4.9	4.7
Conduct disorder (CD)	Moderate	Moderate	50.2	66.8	5.4	4.6
Substance Use Disorder (past 30 days)	Low risk		45.1		6.4	
PTSD	No evidence of PTSD		32.2		4.9	

Interpretation guidance: the higher of the two severity and category scores (bolded above) should generally be used.

Module	Diagnosis
Child/Substance Use Disorder (past 30 days)	None

Diagnostic Probabilities

Module	Probability Measure	Value
Suicide	Probability of Suicide Attempt (next 3 months)	0.024
Anxiety	Probability of Generalized Anxiety Disorder (GAD)	0.058
Mania	Probability of Bipolar Disorder (BP)	0.027
ODD	Probability of Oppositional Defiant Disorder (ODD)	0.964
ADHD	Probability of Attention Deficit Hyperactivity Disorder (ADHD)	0.903
Depression	Probability of Major Depressive Disorder (MDD)	0.020
CD	Probability of Conduct Disorder (CD)	0.755

Note. If the Probability of Suicide Attempt (next 3 months) is .10 or greater, the Suicide row will be displayed in red ink.

Suicide

Timeframe: Past 2 weeks

Category: Moderate

Severity: 47.2

Precision: 4.6

Probability of Suicide Attempt (next 3 months): 0.024

Item Details – Child

Item text	Response	Duration	Severity	Category	Precision
I felt alone.	Somewhat	-	47.0	Moderate	10.3
I felt worthless (like I don't have value).	Just a little	-	45.6	Moderate	9.1
I stopped enjoying anything.	Just a little	-	49.0	Moderate	5.6
I felt everyone would be better off without me.	Just a little	-	49.5	Moderate	4.8
My life wasn't worth living.	Not at all	-	48.5	Moderate	4.8
I made a suicide attempt (I tried to kill myself).	No	2.7	48.1	Moderate	4.6
I thought about killing myself.	Not at all	2.6	47.4	Moderate	4.6
I had a plan for how I would kill myself.	Not at all	2.4	47.2	Moderate	4.6

Anxiety

Timeframe: Past 2 weeks

Child	Parent
Category: Mild Severity: 34.0 Precision: 6.4	Category: Normal Severity: 18.1 Precision: 5.0
Probability of GAD: 0.058	

Item Details – Child

Item text	Response	Duration	Severity	Category	Precision
I felt uneasy much of the time.	Just a little	6.6	43.3	Mild	9.1
Once I started worrying, I couldn't stop.	Not at all	2.4	36.1	Mild	8.0
I had many worries.	Not at all	4.2	34.1	Mild	7.9
I worried about things.	Just a little	3.2	34.7	Mild	7.4
It was hard for me to relax.	Just a little	3.0	34.5	Mild	7.1
My worries overwhelmed me.	Not at all	3.1	33.3	Mild	6.9
I was a worry wart (always worried, nervous).	Not at all	2.8	33.1	Mild	6.9
I worried about how well I did everything.	Just a little	4.2	33.9	Mild	6.6
I could not keep myself from worrying about things.	Not at all	3.1	33.7	Mild	6.6
I just couldn't relax.	Just a little	2.5	34.0	Mild	6.4

Item Details – Parent

Item text	Response	Duration	Severity	Category	Precision
My child worried a lot.	Just a little	7.0	36.8	Mild	11.4
My child worried about things.	Just a little	2.5	34.8	Mild	10.9
My child felt worried.	Just a little	3.0	34.7	Mild	10.4
My child was shy.	Not at all	3.3	23.5	Normal	7.4
My child felt shy around people they were not familiar with.	Not at all	3.1	18.2	Normal	6.6
My child felt nervous.	Just a little	2.1	18.9	Normal	6.4
My child worried too much.	Just a little	2.4	19.4	Normal	6.3
My child worried more than they needed to.	Just a little	3.7	20.2	Normal	6.1
My child was a worrier.	Not at all	2.5	18.8	Normal	6.2
My child worried about how well they do things.	Just a little	3.1	18.8	Normal	6.1
My child worried about other people liking them.	Somewhat	3.3	21.0	Normal	5.6
My child was nervous in new situations.	Not at all	2.7	18.0	Normal	5.5
My child felt afraid that they would make a fool of themselves in front of other people.	Not at all	3.9	16.4	Normal	5.5
My child avoided situations that made them nervous.	Not at all	2.9	15.0	Normal	5.4
My child had trouble relaxing.	Somewhat	4.7	18.1	Normal	5.0

Mania

Timeframe: Past 2 weeks

Child	Parent
Category: Mild Severity: 32.1 Precision: 5.5	Category: Moderate Severity: 60.3 Precision: 4.6
Probability of BP: 0.027	

Item Details – Child

Item text	Response	Duration	Severity	Category	Precision
I was much more interested in sex than usual.	Just a little	5.8	40.7	Mild	14.6
I had times of extreme happiness and energy when some of the things I saw, heard, smelled, tasted, or touched seemed especially vivid (clear, real) or intense.	Not at all	4.8	27.1	Normal	11.2
I had sudden changes in mood or feelings.	Just a little	3.6	28.2	Normal	8.7
I felt excited and happy for no clear reason.	Not at all	3.0	22.8	Normal	8.1
I was more focused on sex than usual (e.g., talking dirty, exposing myself, playing with my private parts, masturbating, making sex phone calls, playing sex games, or touching others sexually).	Just a little	3.1	23.1	Normal	8.1
At times I was much more talkative than usual.	Somewhat	4.9	32.1	Mild	5.5

Item Details – Parent

Item text	Response	Duration	Severity	Category	Precision
My child was physically restless, unable to sit still, and had to keep moving from one activity to another.	Quite a bit	3.0	59.8	Moderate	8.6
For a change, my child found it very exciting to get involved in dangerous, risky, challenging, or emotionally intense activities.	Quite a bit	9.6	68.2	Moderate	7.3
My child frequently felt speeded-up.	Just a little	5.6	63.9	Moderate	6.1
Unlike what is normal for them, my child's attention kept jumping from one idea to another.	Somewhat	4.8	63.2	Moderate	5.2
When my child was talking, it was difficult to interrupt them.	Somewhat	4.8	62.8	Moderate	4.6
Unlike normal, my child's thoughts jumped from topic to topic.	Just a little	4.1	60.3	Moderate	4.6

Oppositional defiant disorder (ODD)

Timeframe: Past 2 weeks

Child	Parent
Category: Severe Severity: 66.7 Precision: 4.3	Category: Severe Severity: 80.1 Precision: 4.2
Probability of ODD: 0.964	

Item Details – Child

Item text	Response	Duration	Severity	Category	Precision
I was rude to adults.	Quite a bit	3.2	60.1	Moderate	7.9
I was bad-tempered.	Somewhat	4.9	58.3	Moderate	6.6
I told an adult I would not follow a rule I didn't like.	Quite a bit	3.6	63.2	Moderate	5.6
I was a hothead.	Very much	4.1	67.0	Severe	5.2
I was argumentative (often argued with others).	Quite a bit	3.0	66.5	Severe	4.7
I was angry because an adult told me what to do.	Quite a bit	4.3	66.7	Severe	4.3

Item Details – Parent

Item text	Response	Duration	Severity	Category	Precision
My child was easy to get along with.	Not at all	5.3	83.7	Severe	11.3
My child was rude to adults.	Quite a bit	2.4	80.6	Severe	6.9
My child was defiant with adults.	Very much	2.6	84.3	Severe	5.9
My child got angry frequently.	Quite a bit	2.8	81.8	Severe	4.9
My child argued with a parent, teacher, or boss.	Quite a bit	2.9	79.0	Moderate	4.5
My child was like a bomb that was ready to explode.	Quite a bit	3.7	80.1	Severe	4.2

Attention deficit/hyperactivity disorder (ADHD)

Timeframe: Past 2 weeks

Child	Parent
Category: Moderate Severity: 57.9 Precision: 6.0	Category: Moderate Severity: 62.0 Precision: 3.4
Probability of ADHD: 0.903	

Item Details – Child

Item text	Response	Duration	Severity	Category	Precision
I had trouble staying focused.	Somewhat	3.6	43.3	Mild	11.4
I was easily distracted by unimportant things when I was trying to do my schoolwork or other work.	Somewhat	4.1	42.3	Mild	10.7
I was easily distracted.	Somewhat	2.5	43.1	Mild	10.3
I forgot to do something that I do regularly, like going to a meeting or a practice.	Somewhat	3.6	47.6	Mild	9.2
I demanded a lot of attention from other people.	Somewhat	4.4	51.5	Moderate	7.9
I interrupted other people when they were speaking.	Quite a bit	4.8	53.5	Moderate	7.7
I moved from one task to another without completing any of them.	Somewhat	3.8	55.5	Moderate	7.3
My attention wandered from one thing to another.	Quite a bit	4.0	55.5	Moderate	7.3
I had trouble paying attention to important things for more than a short time.	Quite a bit	2.6	56.4	Moderate	7.2
I had a hard time settling down after an exciting activity.	Somewhat	3.8	57.9	Moderate	6.0

Item Details – Parent

Item text	Response	Duration	Severity	Category	Precision
My child liked to play so wild and recklessly that they could get hurt.	Quite a bit	3.7	65.9	Moderate	10.6
My child had difficulty playing quietly.	Somewhat	3.0	63.8	Moderate	7.1
My child had trouble sitting still when they were supposed to (at movies, church, etc.).	Somewhat	3.4	62.5	Moderate	5.9
My child left their seat when they weren't allowed to get up.	Somewhat	3.8	61.6	Moderate	5.1
My child seemed restless when they had to sit for more than ten minutes.	Quite a bit	3.6	63.0	Moderate	4.6
My child got up and left their seat in school or other situations when they were supposed to stay seated.	Somewhat	4.5	62.0	Moderate	3.4

Depression

Timeframe: Past 2 weeks

Child	Parent
Category: Mild Severity: 48.9 Precision: 4.9	Category: Mild Severity: 40.4 Precision: 4.7
Probability of MDD: 0.020	

Item Details – Child

Item text	Response	Duration	Severity	Category	Precision
I felt everyone would be better off without me.	Just a little	4.4	49.3	Mild	9.7
My life wasn't worth living.	Not at all	4.5	46.0	Mild	9.1
I stopped enjoying anything.	Just a little	2.3	50.1	Moderate	6.1
I felt worthless (like I don't have value).	Just a little	2.7	50.0	Moderate	5.8
I felt alone.	Somewhat	3.4	50.6	Moderate	5.7
I was less motivated to do things than usual (I felt there was no longer a reason to do things).	Just a little	3.3	48.9	Mild	4.9

Item Details – Parent

Item text	Response	Duration	Severity	Category	Precision
My child seemed to stop enjoying anything.	Somewhat	3.5	55.5	Moderate	6.4
My child thought life wasn't worth living.	Not at all	5.2	53.1	Moderate	6.0
Even little things seemed to make my child more tired than usual.	Not at all	5.0	47.9	Mild	5.3
My child felt down in the dumps.	Just a little	5.0	46.4	Mild	5.2
My child didn't have fun doing anything.	Not at all	7.5	43.7	Mild	4.8
My child was more quiet than usual.	Not at all	2.8	40.4	Mild	4.7

Conduct disorder (CD)

Timeframe: Past 2 weeks

Child	Parent
Category: Moderate Severity: 50.2 Precision: 5.4	Category: Moderate Severity: 66.8 Precision: 4.6
Probability of CD: 0.755	

Item Details – Child

Item text	Response	Duration	Severity	Category	Precision
I got so mad that I broke things.	Somewhat	3.4	40.0	Mild	13.2
I fought a lot with others my own age.	Quite a bit	3.1	50.3	Moderate	10.5
I used a weapon that could cause serious physical harm to others (e.g., a bat, brick, broken bottle, knife, or gun).	No	4.5	42.2	Moderate	8.0
I got into physical fights when I was upset.	Somewhat	2.7	43.9	Moderate	7.1
I tried to cheat people by selling or trading things that weren't what I said they were.	Just a little	9.6	50.1	Moderate	5.4
I hit another person when I was provoked.	Somewhat	3.9	50.2	Moderate	5.4

Item Details – Parent

Item text	Response	Duration	Severity	Category	Precision
My child stole from home, school, or elsewhere.	Somewhat	3.0	54.9	Moderate	10.5
My child stole things when they thought no one was watching.	Somewhat	3.2	54.6	Moderate	10.5
My child destroyed things on purpose.	Somewhat	3.2	60.7	Moderate	6.2
My child used force, or threatened to use force, to make other people do what they wanted.	Quite a bit	3.5	66.8	Moderate	5.0
My child threatened people they know.	Quite a bit	2.2	66.7	Moderate	4.7
My child picked on other kids.	Quite a bit	2.5	66.8	Moderate	4.6

Substance Use Disorder (past 30 days)

Part 1: Substance Use and Diagnosis

Timeframe: Past 30 days

SUD Diagnosis:

None

Item Details

Item text	Response	Duration	Severity	Category	Precision
Have you been incarcerated in the past 3 months?	No	2.8	-		-
Have you been hospitalized in a drug treatment center in the past 3 months?	No	1.4	-		-
Have you been on probation in the last 3 months?	No	1.1	-		-
Have you used any of the following in the last month, other than prescriptions from your doctor (select one or more of the following or "None of these"):	Alcohol	10.0	-		-
Have you used any of the following in the last month, other than prescriptions from your doctor (select one or more of the following or "None of these"):	None of these	2.0	-		-
Have you used any of the following in the last month, other than prescriptions from your doctor (select one or more of the following or "None of these"):	None of these	2.0	-		-
When I stop using alcohol I get withdrawal symptoms (for example, sweating, heart racing, shakiness, seizures, nausea/vomiting, hallucinations).	No	2.4	-		-
In the past month have you spent a lot of time using, obtaining, thinking about or craving alcohol?	No	2.8	-		-

Item text	Response	Duration	Severity	Category	Precision
In the past month have you or someone else wanted you to cut back on alcohol?	No	3.6	-		-
In the past month has alcohol caused problems for you or kept you from getting things done?	Yes	4.6	-		-
In the past month, have you used alcohol again and again, despite the harm it has caused?	No	2.7	-		-

Substance Use Disorder (past 30 days)

Part 2: Current Severity, Category and Precision

Timeframe: Past 30 days

Category: Low risk

Severity: 45.1

Precision: 6.4

Item Details

Item text	Response	Duration	Severity	Category	Precision
How troubled or bothered have you been in the past 30 days by alcohol problems?	Slightly	7.5	57.2	Intermediate risk	13.1
Have you been preoccupied with using alcohol and/or using other drugs like marijuana, or with taking medications either without a prescription or more than they were prescribed?	No	2.8	48.2	Low risk	11.3
How troubled or bothered have you been in the past 30 days by drug problems?	Not at all	3.4	47.3	Low risk	11.0
How much were you bothered by feelings of guilt?	Moderately	4.0	52.1	Intermediate risk	8.8
How much were you bothered by feelings of worthlessness?	A little bit	2.8	52.8	Intermediate risk	8.5
How much were you bothered by having strong negative feelings such as fear, horror, anger, guilt, or shame?	Not at all	3.1	49.3	Low risk	8.4
How much were you bothered by blaming yourself for things?	A little bit	7.5	48.9	Low risk	8.4
Did you need to take sedatives, tranquilizers, sleeping pills or muscular relaxants to deal with the problems in your life (examples: Xanax, Ativan, Valium, Ambien, Lunesta, Flexeril, Robaxin)?	Never	2.5	46.4	Low risk	7.5

Item text	Response	Duration	Severity	Category	Precision
How much were you bothered by feeling blue (down or sad)?	A little bit	3.6	46.7	Low risk	7.4
How much have you been emotionally affected by your health condition?	Mild	3.1	47.2	Low risk	6.9
How much were you bothered by feeling very upset when something reminded you of a stressful experience from the past?	Not at all	4.3	46.1	Low risk	6.9
How much were you bothered by feeling jumpy or easily startled (surprised)?	A little bit	5.1	46.5	Low risk	6.9
Please rate how much your problems get in the way of your ability to carry out certain activities, where 1 is not at all and 9 is severely. Because of my [problem], I have difficulty doing my work.	3	3.4	46.3	Low risk	6.8
Have you taken sedatives, hypnotics, benzodiazepines, minor tranquilizers, or muscular relaxants (examples: Xanax, Ativan, Valium, Ambien, Lunesta, Flexeril, Robaxin)?	No	2.1	44.4	Low risk	6.5
How much were you bothered by a loss of interest in activities that you used to enjoy?	Moderately	4.2	45.1	Low risk	6.4

PTSD

Timeframe: Past 2 weeks

Category: No evidence of PTSD

Severity: 32.2

Precision: 4.9

Item Details

Item text	Response	Duration	Severity	Category	Precision
How much were you bothered by trying to stay away from anything that reminds you of a scary or very upsetting thing? (For example, people, places, or conversations about the scary or very upsetting thing)	Not at all	4.0	23.1	No evidence of PTSD	11.8
How much were you bothered by trying not to think about or have feelings about a scary or very upsetting thing that happened?	Not at all	4.8	17.3	No evidence of PTSD	10.9
How much were you bothered by having strong bad feelings (like fear, anger, guilt, or shame)?	Moderately	4.5	34.6	No evidence of PTSD	6.7
How much were you bothered by being jumpy or easily scared? (For example, when someone walks up behind you, when you hear a loud noise)	Not at all	4.1	30.7	No evidence of PTSD	6.1
How much were you bothered by feeling upset when you remembered a scary or very upsetting thing? (For example, feeling scared, angry, sad, guilty, confused)	A little bit	6.1	33.6	No evidence of PTSD	5.1
How much were you bothered by acting or feeling as if a scary or very upsetting thing was happening again? (For example, seeing or hearing something and feeling as if you are in	Not at all	3.6	32.2	No evidence of PTSD	4.9

Item text	Response	Duration	Severity	Category	Precision
the scary or very upsetting thing again)					

End of Report